



a 2 - œ-fp

- 1! 3 « -TMµ j 3 © TMµ -@Y TMk TMk - qýx - 3 3 » 3 © - « » © » j © - qýx - µ @k a l ° - » j vQ
- 3 « 2 j ° » » © qýx - 3 Q
- 2! 3 « -@j | ĩ ñ œ « 1 © » j w © -@j | q © « | ĩ « 1 © ĩ ñ œ - 2 2 ° - | © - ° - vQ
- 3! 3 « ĩ n « 1 © » j 3 © q - 2 2 ° µ o - © - ° µ TM± TM 10-15 2 - © - vQ
- 4! 3 « ° ° ĩ mō « TM » © TM - , 2 µ j 3 Q

@ - , á © ©

° © - 2 - m - 3 TM - @ 3 » - , - @ - , á © - » » Q

. 1 - -

-p - , x . 1 ž - , - » » Q µ ¼ - ° ĩ o µ j - Q

- µ j | ø - - @x

- 1 - x - 2 | { - - y 3 » - j ° 3 © - © | - ĩ n ° « ĩ j 2 - 2 TM È « - œ » vQ
- 2 3 TM - 3 TM » | ø - « ° ĩ q © TM - µ w ° » ĩ µ - Y j Q
- 3 | ø ž 3 ° œ » p - x - ž © 3 - ĩ - - - @x ĩ o - - ĩ - ĩ j ° ĩ n ° © - ĩ n ° « ĩ - - - © - 1 µ - » Q

° » - , ©

- 1 - x - , - @ - ĩ - µ µ TM 2 2 ĩ 2 ĩ m a . ĩ TM 2 ĩ « @k a l ĩ j - µ - ĩ m a . ĩ TM 2 ĩ µ µ TM µ µ TM µ µ 2 ĩ 2 © µ ĩ œ ° - ĩ m « ĩ j - œ » 3 Q
- 2 j ° - 2 ĩ m a . ĩ TM 2 ĩ - x - , - ± ð ĩ - - » » m ç Y © » - » n ž ° - ž . « ĩ © - , 3 á © - ĩ µ ú ° ĩ œ | » - œ » 3 ĩ ° - x - , - ± © . © ĩ « 1 © - » - © 3 Q

° 3 - - œ -

- µ ĩ 3 © ĩ	#!&# ĩ ± ĩ ± ĩ
ĩ 2 - ĩ p	#!%# ĩ ± ĩ ± ĩ
° oö ĩ ĩ 3 2 ĩ µ ä - ĩ ° ý - - x /	(!# ĩ ± ĩ ± ĩ
- ĩ 3 ĩ ĩ © ĩ	@3 b © 2 ĩ ĩ
	3 TM - TSSISS ĩ ± ĩ ± ĩ

- 2 | ©

m j 2 ĩ µ - TM ĩ m - °

qñ © « - © -

m j 2 ĩ µ - TM ĩ m - °

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

# FIPGOLD

**Fipronil 0.3% G.R.**

( Insecticide )

Fipronil 0.3% GR is a insecticide containing Fipronil 0.3% for control of rice pest complex and Early shoot borer & Root borer in Sugarcane and termite in wheat.

## **Recommendation**

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water	Waiting Period between last spray to harvest	Re-entry after each Application (In Hours)
		AI (gm)	Formulation (kg)			
Sugarcane	Root borer, Early shoot borer	75-100	25.0-33.3		9 months	
Wheat	Termites	0.06	20kg	-----	91	
Rice	Green leaf hopper, Brown plant hopper, Stem borer, Rice Leaf Folder, Rice Gall Midge, White backed Plant hopper, Whorl maggot	50-75	16.67-25.00		32 days	

## **Direction of Use**

Method of Application and Equipment for use : Recommended dosage for granule should be broadcasted using operated granule applicator or mechanical dispenser application in Paddy, 2-3 cm standing water should be in the field and keep water impounded for 2-3 days after application.

## **Precaution**

1. Keep away from foodstuffs, empty foodstuff containers and animals food.
2. Avoid contact with mouth, eyes and skin.
3. Avoid inhalation while in using.
4. Wash thoroughly the contaminated clothes and parts of the body after broadcasting.
5. Do not smoke, drink, eat and chew anything while broadcasting.
6. Wear full protective clothing while broadcasting.

Warning :- Fipronil is toxic to aquatic animals & honey bee hence avoid the use near aquatic environment & during to rassing period of honey bee

## **Symptoms Of Poisoning**

Irritability, lethargy, muscle tremors , convulsion may occur.

## **First Aid**

1. If swallowed, induce vomiting by tickling the back of throat. Repeat it until the vomitus is clear. Do not induce vomiting if the patient is unconscious.
2. If clothing and skin are contaminated, remove the cloths and wash the contaminated skin with copious amount of soap and water.
3. If eyes are contaminated, flush with plenty of saline/clean water for about 10 to 15 minutes.
4. If inhaled, remove the patient to fresh air.

## **Phytotoxicity**

The product is not phytotoxic when used as per field recommendation.

**Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet**

